

T-odd functions in relativistic eikonal models

Marc Schlegel

Theory Center
Jefferson Lab

in collaboration with Leonard Gumberg

Berkeley Summer Program on Nuclear Spin Physics, June 10, 2009


Generalizations of collinear Parton Distributions

GPDs (off-diagonal):

measurable in exclusive processes
provides 3-dim. spatial picture

TMDs (small transv. deviations):

measurable in semi-inclusive processes
provides 3-dim. momentum picture



$$\Phi_{ij}(x) = \int_{-\infty}^{\infty} \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle P | \bar{q}_j(-\frac{z^-}{2}) [-\frac{z^-}{2}; \frac{z^-}{2}] q_i(\frac{z^-}{2}) | P \rangle$$


Collinear PDFs $f_1(x, \mu), g_1(x, \mu), h_1(x, \mu)$

Theory well-understood, delivers a one-dimensional picture of nucleon structure

Measurable in DIS (but not sensitive to transversity...)

Generalized Parton Distributions

- Exclusive processes (DVCS, meson production, ...):



$$P = \frac{1}{2}(p + p')$$

$$\Delta = p' - p$$

$$\Delta^+ = -2\xi P^+$$

$$t = -\frac{4\xi^2 M^2 + \vec{\Delta}_T^2}{1 - \xi^2}$$

$$F_{ij}(x, \xi, \vec{\Delta}_T) = \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle p' | \bar{q}_j \left(-\frac{z^-}{2} \right) \left[-\frac{z^-}{2}; \frac{z^-}{2} \right] q_i \left(\frac{z^-}{2} \right) | p \rangle$$

Eight GPDs → most prominent H and E


$$P^+ \text{Tr}[F \gamma^+] = \bar{u}(p') \left[\gamma^+ H(x, \xi, t) + \frac{i\sigma^{+\mu} \Delta_\mu}{2M} E(x, \xi, t) \right] u(p)$$

E → no correspondence in the collinear limit. Ji-Angular momentum sum rule

$$J^q = \frac{1}{2} \int_{-1}^1 dx x [H^q(x, 0, 0) + E^q(x, 0, 0)]$$

Impact Parameter Space

- Impact Parameter Space: ($\xi=0$) [M. Burkardt, PRD62, 071503]



- Impact parameter b_T and transv. momentum transfer $\Delta_T \rightarrow FT$

$$\mathcal{F}_{ij}(x, \vec{b}_T) = \int \frac{d^2 \Delta_T}{(2\pi)^2} e^{-i \vec{\Delta}_T \cdot \vec{b}_T} F_{ij}(x, 0, \vec{\Delta}_T)$$



- Impact parameter space → “diagonal” matrix element $z_{1/2} = \mp \frac{z^-}{2} n_- + b_T$

$$\mathcal{F}_{ij}(x, \vec{b}_T) = \int \frac{dz^-}{2(2\pi)} e^{ixP^+ z^-} \langle P^+; \vec{0}_T | \bar{\psi}_j(z_1) [z_1; z_2] \psi_i(z_2) | P^+; \vec{0}_T \rangle$$


- ($\xi=0$): probability density of partons in transverse plane.

k_T -dependent Parton Distributions (TMDs)

Semi-inclusive processes at small transverse final state momenta:



$$\Phi_{ij}(x, \vec{k}_T; S) = \int \frac{dz^- d^2 z_T}{2(2\pi)^3} e^{ik \cdot z} \langle P, S | \bar{\psi}_j(0) \mathcal{W}_{\text{SIDIS/DY}}[0, z] \psi_i(z) | P, S \rangle \Big|_{z^+ = 0}$$



- Eight TMDs:


		quark pol.		
		U	L	T
nucleon pol.	U	f_1		h_1^\perp
	L		g_1	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}	h_1, h_{1T}^\perp

Twist-2 TMDs


$$\frac{1}{2} \text{Tr}[\Phi \gamma^+] = f_1(x, \vec{k}_T^2) - \frac{\varepsilon_T^{ij} k_T^i S_T^j}{M} f_{1T}^\perp(x, \vec{k}_T^2)$$

T-odd TMDs


- **Neglect** gauge link operator:



- **If T-odd TMDs $\neq 0$:** Gauge Link not negligible, **physical effect**:



$$\mathcal{W}[z_1; z_2] = \mathcal{P} e^{-ig \int_{z_1}^{z_2} ds \cdot A(s)}$$



Initial / Final state interactions

Time reversal \longrightarrow switches sign:


$$f_{1T}^\perp \Big|_{DIS} = - f_{1T}^\perp \Big|_{DY}$$

$$h_1^\perp \Big|_{DIS} = - h_1^\perp \Big|_{DY}$$

Illustrative picture of the Final State Interactions

Situation in SIDIS:

Unpolarized nucleon → quarks are equally distributed in transverse plane




Final State interactions are assumed to be attractive here



Lensing effect, but no observable net effect!

Sivers asymmetry

Transversely polarized nucleon →
spatial distortion of the quark distribution in the transverse plane →
Impact parameter GPD E



Spatial distortion + FSI lead to observable net effect
→ non-zero Left-Right (Sivers) asymmetry
→ attractive FSI: correct prediction of the sign of SSA

Relation Sivers – E

Non-trivial relations for “T-odd” parton distributions:

M. Burkardt [Nucl.Phys. A735, 185], [PRD66, 114005]

Average transverse momentum of unpolarized partons
in a transversely polarized nucleon:

$$\langle k_T^i \rangle_T(x) = \int d^2 k_T k_T^i \frac{1}{2} \left[\text{Tr}[\gamma^+ \Phi(\vec{S}_T)] - \text{Tr}[\gamma^+ \Phi(-\vec{S}_T)] \right] \propto f_{1T}^{\perp,(1)}(x)$$



Manipulation of Gauge Links + Impact parameter representation

$$\langle k_T^i \rangle(x) = \int d^2 b_T \int \frac{dz^-}{2(2\pi)} e^{ixP^+ z^-} \langle P^+; \vec{0}_T; S_T | \bar{\psi}(z_1) \gamma^+ [z_1; z_2] I^i(z_2) \psi(z_2) | P^+; \vec{0}_T; S_T \rangle$$

$$z_{1/2} = \mp \frac{z^-}{2} n_- + b_T$$

Impact parameter representation for GPD E

$$I^i(z^-) = \int dy^- [z^-; y^-] gF^{+i}(y^-) [y^-; z^-] \text{ coll. “soft gluon pole” matrix element}$$


- **(Semi-) classical approximation:** $\hat{I}^i \simeq \mathcal{I}^i(x, \vec{b}_T) \mathbf{1} + \dots$
 → factorization of **final state interactions** and **spatial distortion**:

$$\langle k_T^i \rangle = -M \epsilon_T^{ij} S_T^j f_{1T}^{\perp, (1)}(x) \simeq \int d^2 b_T \mathcal{I}^i(x, \vec{b}_T) \frac{\vec{b}_T \times \vec{S}_T}{M} \frac{\partial}{\partial b_T^2} \mathcal{E}(x, \vec{b}_T^2)$$


$\mathcal{I}^i(x, \vec{b}_T^2)$: Lensing Function = net transverse momentum

- Also for Boer-Mulders function and chirally-odd GPDs...
- Relation valid for lowest order diagrams in simple spectator models:
 [Burkardt, Hwang, PRD69, 074032], [Meissner, Metz, Goeke, PRD76, 034002]

GPDs:




(T-odd) TMDs:



- Analysis of *mother functions*: No exact, model-independent relations possible.
 [Meissner, Metz, MS, Goeke, JHEP08(2008)038]
- "How approximate" is the relation? Need to know the lensing function better...


FSI beyond One-Gluon exchange

- Final State Interactions (Sivers-effect) mostly modeled by a One-Gluon Exchange:





Example Diquark-spectator:
Sivers function $\propto \alpha_s$
Sivers effect at HERMES $\approx 5\%$
 $\Rightarrow \alpha_s \approx 0.2 - 0.3$


- Relativistic Eikonal models: Treat FSI non-perturbatively.



- Work in the diquark picture (Lens. funct. identifiable)
→ more realistic for a pion...
- Only diagrams that reflect the "naive picture".



Lensing Function



Assume a non-perturbative scattering amplitude $M +$
Separate GPD and FSI via contour integration

$$\int \frac{d^4 q}{(2\pi)^4} g_N [(P - q)^2] \frac{[(\not{P} - \not{q} + m_q) u(P, S)]_i \mathcal{M}_{bc}^{ab}(q, P - k)}{[n \cdot (P - k - q) + i\varepsilon][(P - q)^2 + m_q^2 + i\varepsilon][q^2 - m_s^2 + i\varepsilon]}$$

- **Step 1: Integration over q^- :**

Assume no q^- & q^+ poles in M .

q^- - poles at one loop for higher twist T-odd TMDs [Gamberg, Hwang, Metz, MS, PLB 639, 508]
→ non-light like Wilson lines, would spoil relation.

- **Step 2: Integration over q^+ :**

Fix the q^+ - pole

emphasizes a "natural" picture of FSI

equivalent to Cutkosky cut, assumptions of Step 1 valid in Eikonal models

$$\frac{1}{(1-x)P^+ - q^+ + i\varepsilon} = P \frac{1}{(1-x)P^+ - q^+} - i\pi\delta((1-x)P^+ - q^+)$$


$$I^i(x, \vec{q}_T) = \int \frac{d^2 p_T}{(2\pi)^2} (2p_T - q_T)^i \Im M_{bc}^{ab}(|\vec{p}_T|) \left((2\pi)^2 \delta^{ac} \delta^{(2)}(\vec{p}_T - \vec{q}_T) + \Re M_{da}^{cd}(|p_T - q_T|) \right)$$

Relativistic Eikonal Models

Scattering amplitude of two highly energetic particles at small momentum transfer:

Possible under two assumptions:

1) Particles are eikonalized (not even an appr. for quark...) 2) Gener. Ladder Approx.



In QED: Coulomb phase (70s)

- 1) Diagrammatic: Wu, Yang,...
- 2) Combinatorial: Levy, Sucher, Quiros...
- 3) Functional: Fried, Gabellini,...

Here: Outline of the functional approach for quark-antiquark scattering (pion):

[Fried, Gabellini, Avan, Eur.Phys.J. C13, 699]

QCD-generating functional:

$$Z[j, \eta, \bar{\eta}, \xi, \bar{\xi}] = \mathcal{N} \int \mathcal{D}A e^{-\frac{i}{4} \int F^2 - \frac{i}{2} \int (\partial \cdot A)^2 + i \int j \cdot A} e^{\text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} e^{-i \int \bar{\eta} G[A] \eta - i \int \bar{\xi} H[A] \xi}$$

Quark-Antiquark 4-point function:

$$\begin{aligned} T(i_1, i_2 | f_1, f_2) &= \frac{\delta}{\delta \bar{\eta}(f_1)} \frac{\delta}{\delta \eta(i_1)} \frac{\delta}{\delta \bar{\eta}(i_2)} \frac{\delta}{\delta \eta(f_2)} Z \Big|_{j=\eta=\bar{\eta}=\xi=\bar{\xi}=0} \\ &= -\mathcal{N} \int \mathcal{D}A e^{-\frac{i}{4} \int F^2 - \frac{i}{2} \int (\partial \cdot A)^2} e^{\text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} G(i_2, f_2 | A) G(f_1, i_1 | A) \end{aligned}$$

Eikonal Approximation

Idea: highly energetic particle loses spin information $\gamma^\mu \longrightarrow v^\mu$

In QED:


$$\begin{aligned} G(x, y|A) &= (i\partial - m_q + A(x))^{-1} \delta^{(4)}(x - y) \\ &\longrightarrow (v \cdot (i\partial + A(x)) - m_q)^{-1} \delta^{(4)}(x - y) = -i \int_0^\infty ds e^{is(v \cdot (\partial + A(x)) - m_q + i\epsilon)} \delta^{(4)}(x - y) \end{aligned}$$

$$G_{\text{eik}}^{ab}(x, y|A) = -i \int_0^\infty ds e^{-is(m_q - i0)} \delta^{(4)}(x - y - sv) \left(e^{-ig \int_0^s d\beta v \cdot A^\alpha(y + \beta v) t^\alpha} \right)_+^{ab}$$

Trick to disentangle the A-field and the color matrices t: Functional FT

$$\left(e^{-ig \int_0^s d\beta v \cdot A^\alpha(y + \beta v) t^\alpha} \right)_+^{ab} = \mathcal{N}' \int \mathcal{D}\alpha \int \mathcal{D}u e^{i \int d\tau \alpha^\beta(\tau) u^\beta(\tau)} e^{ig \int d\tau \alpha^\beta(\tau) v \cdot A^\beta(y + \tau v)} \left(e^{i \int_0^s d\tau t^\beta u^\beta(\tau)} \right)_+^{ab}$$

Transformation of the scattering amplitude into momentum space + Amputation
+ Insertion of eikonal propagators:



$$\left(\frac{\partial}{\partial g} G^{\text{eik}} \right)_{\text{amp}} (P_2, P_1)$$

Eikonal Amplitude

$$\frac{\partial^2 M}{\partial g_1 \partial g_2} \simeq \int \mathcal{D}A e^{i \int [-\frac{1}{4} F^2 - \frac{\lambda}{2} (\partial \cdot A)^2] + \text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} \left(\frac{\partial G}{\partial g_1} \right) \left(\frac{\partial \bar{G}}{\partial g_2} \right)$$

Further steps (in words) [Fried, Gabellini, Avan, Eur.Phys.J. C13, 699]:

- Use scale invariance of the gauge vector n : "Quasi-Abelian limit"
- Separate off quadratic gluon terms
- Neglect Fermion loops and gluon self-interactions (Ladder approximation)

Final result for the eikonal quark-antiquark scattering amplitude:

$$M_{ab}^{\text{eik}}(q_T) = \frac{2(1-x)P^+}{m_s} \int d^2 z_T e^{-iq_T \cdot z_T} \left[\int d^{N^2-1} \alpha \int \frac{d^{N^2-1} u}{(2\pi)^{N^2-1}} (e^{i\chi(|z_T|)t \cdot \alpha})_{ac} (e^{it \cdot u})_{cb} e^{-i\alpha \cdot u} - 1 \right]$$

Abelian U(1)-theory: recover Coulomb phase

$$\dots \left[e^{i\chi(|z_T|)} - 1 \right]$$

SU(2): analytical evaluation of color effects:

$$e^{i\frac{\sigma}{2} \cdot u} = \cos\left(\frac{|u|}{2}\right) + i \frac{\sigma \cdot u}{|u|} \sin\left(\frac{|u|}{2}\right)$$

SU(3): (?) numerical evaluation, in process...

The Eikonal Phase

Recall: $\chi(|z_T|) = g^2 \int_{-\infty}^{\infty} d\tau_1 \int_{-\infty}^{\infty} d\tau_2 n^\mu \bar{n}^\nu \mathcal{D}_{\mu\nu}(z + \tau_1 n - \tau_2 \bar{n}) = \frac{\alpha}{\pi} \int d^2 k_T n^\mu \bar{n}^\nu \tilde{\mathcal{D}}_{\mu\nu}(k_T) e^{i\vec{k}_T \cdot \vec{z}_T}$

For a free gluon propagator: $\chi(|z_T|) = 2\alpha K_0(\mu|z_T|)$


- **Good news:** Recover exactly old perturbative results in the one gluon exchange.
[Burkardt, Hwang; Meißner, Metz, Goeke]

e.g.: scalar diquark: $\mathcal{I}^i(x, \vec{b}_T) = -\alpha(1-x)\frac{b_T^i}{\vec{b}_T^2}$ $f_{1T}^\perp(x, \vec{k}_T^2) = -\frac{g^2}{2(2\pi)^3}\alpha \frac{(1-x)M(xM+m_q)}{\vec{k}_T^2(\vec{k}_T^2+\tilde{m}^2)} \ln\left(\frac{\vec{k}_T^2+\tilde{m}^2}{\tilde{m}^2}\right)$

- **Bad news:** Dependence on IR-regulator.

Exchanged gluons are soft \rightarrow free propagator? What is the value of α_s in the IR-limit?

→ **Input from Dyson-Schwinger Equations.**



[Plot from Fischer, Maas, Pawłowski, arXiv:0810.1987]

- Calculations in Landau gauge + Euclidean space \rightarrow applicable here.
- IR-finite gluon propagator $\mathcal{D}(p^2 \rightarrow 0) \sim (p^2)^{(2\kappa-1)}$
- Renorm. Point at $\mu_R = m_Z$ \rightarrow here: $\mu_R = 1\text{GeV}$
- DSE: IR-limit of $\alpha_s(p^2 \rightarrow 0) \sim 2.972$
- Use running of α_s as vertex factor.

Lensing Function



Express Lensing Function in terms of Eikonal Phase:

$$\mathcal{I}_{(N=1)}^i(x, \vec{b}_T) = \frac{1}{4} \frac{\vec{b}_T^i}{|\vec{b}_T|} \chi'(\frac{|\vec{b}_T|}{1-x}) \left[1 + \cos \chi(\frac{|\vec{b}_T|}{1-x}) \right]$$

$$\mathcal{I}_{(N=2)}^i(x, \vec{b}_T) = \frac{1}{4} \frac{\vec{b}_T^i}{|\vec{b}_T|} \chi'(\frac{|\vec{b}_T|}{1-x}) \left[1 + \cos \frac{\chi}{4} + \frac{\chi}{8} (\frac{\chi}{4} - \sin \frac{\chi}{4}) \right] (\frac{|\vec{b}_T|}{1-x})$$

$$\mathcal{I}_{(N=3)}^i(x, \vec{b}_T) = ???$$


Eikonal Phase



Final State Interactions remain negative (even in SU(2))



The GPD E in the diquark model

The remaining piece of the puzzle: GPD E



Scalar diquark with a multi-pole diquark vertex factor



$$\Lambda^{2(\alpha-1)} \frac{k^2 - m_q^2}{[k^2 - \Lambda^2]^\alpha}$$



- Need to fit the model parameters:
GPD-limits:
Form Factors F1, F2 → t-dependence
+ valence $u(x)$ (GRV) → x-dependence
- Difficult to fit all limits for diquark only
→ Form Factor fit preferred.
- Other GPD model:
[Diehl et al., EPJ C39 (2005) 1-39]

$$E_v^u(x, 0, t) = (N^u \kappa^u x^{-\alpha} (1-x)^\beta) e^{tg^u(x)}$$

Results for the u -quark Sivers function



Extraction by Torino-Group

$$\Delta^N f^{(1)}(x) = -f_{1T}^\perp(x)$$

- Relation produces a Sivers effect $\sim 0.04 - 0.08$, Torino - Extraction ~ 0.05 .
- SU(2) color factors seem to increase the effect. Effect of SU(3) ???
- Relation overshoots extraction, but correct sign and order of magnitude...

Summary:

- Relation Sivers – E via separation of FSI + spatial distortion of parton dist.
- Relation is not rigorous, model-dependent.
Holds for lowest order spectator models.
- Relativistic Eikonal model: Non-perturbative, field-theoretical model of FSI
→ Lensing Function.
- Relation reproduces the right order of magnitude of the Sivers effect.

Outlook:

- Apply formalism for other soft objects:
Boer-Mulders, Collins function, soft factor...
- Improve the diquark model through axial-vectors.
- SU(3) color factors
- Other Dyson-Schwinger extractions of Gluon-Prop. / running coupling.

The Quasi-Abelian Limit

$$\left(\frac{\partial}{\partial g} G^{\text{eik}} \right)_{\text{amp}}^{ab}(P_2, P_1) = \mathcal{N}' \int d^4 z e^{-iz \cdot (P_1 - P_2)} \int \mathcal{D}\alpha \int \mathcal{D}u \alpha^\beta(0) v \cdot A^\beta(z) e^{ig \int_{-\infty}^{\infty} d\tau \alpha^\beta(\tau) v \cdot A^\beta(z + \tau v)} e^{i \int \alpha \cdot u} \left(e^{i \int_{-\infty}^{\infty} d\tau t^\beta u^\beta(\tau)} \right)_+$$

Description of the gauge link: Invariance under rescaling of gauge vector: $v = n \rightarrow \lambda n$

$$\int_{-\infty}^{\infty} \lambda d\tau \alpha^\beta(\tau) n \cdot A^\beta(z + \tau \lambda n) = \int_{-\infty}^{\infty} d\tilde{\tau} \alpha^\beta\left(\frac{\tilde{\tau}}{\lambda}\right) n \cdot A^\beta(z + \tilde{\tau} n) \simeq \alpha^\beta(0) \int_{-\infty}^{\infty} d\tau n \cdot A^\beta(z + \tau n)$$

"Quasi-Abelian Limit" [Fried, Gabellini, PRD 55,2430]

For spectator eikonal propagator: $\lambda \sim \frac{(1-x)P^+}{m_s}$ **"Infinite momentum frame"**

Back to the eikonal amplitude: Insertion of the quasi-abelian limit.

$$\frac{\partial^2 M}{\partial g_1 \partial g_2} \simeq \int \mathcal{D}A e^{i \int [-\frac{1}{4} F^2 - \frac{\lambda}{2} (\partial \cdot A)^2] + \text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} \left(\frac{\partial G}{\partial g_1} \right) \left(\frac{\partial \bar{G}}{\partial g_2} \right)$$

Eikonal Amplitude

Focus on the gluonic part: Separate off quadratic gluon terms

$$\frac{\partial^2 M}{\partial g_1 \partial g_2} \simeq \int \mathcal{D}A e^{i \int [-\frac{1}{4} F^2 - \frac{\lambda}{2} (\partial \cdot A)^2] + \text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} \left(\frac{\partial G}{\partial g_1} \right) \left(\frac{\partial \bar{G}}{\partial g_2} \right)$$

Write it in terms of a Gaussian integral:

$$\propto \dots \left[\int \mathcal{D}A e^{-\frac{i}{2} \int A \mathcal{D}^{-1} A + i \int J \cdot A} \mathcal{F}[A] \right]$$

$$\mathcal{F}[A] = e^{-\frac{i}{4} \int (F^2 - f^2)} e^{\text{Tr} \ln G^{-1}[A] + \text{Tr} \ln H^{-1}[A]} = \int \mathcal{D}\phi \tilde{\mathcal{F}}[\phi] e^{i \int A \cdot \phi}$$

\mathcal{D}^{-1} : inverse free gluon propagator

Eikonal current in the quasi-abelian limit:

$$J^{\mu, \beta}(x) = g_1 \alpha^\beta(0) \int_{-\infty}^{\infty} d\tau n^\mu \delta^{(4)}(x - z - \tau n) + g_2 \bar{\alpha}^\beta(0) \int_{-\infty}^{\infty} d\tau \bar{n}^\mu \delta^{(4)}(x - \bar{z} + \tau \bar{n})$$

Manipulation via functional translation: $\mathcal{F}[A + B] = e^{\int G \frac{\delta}{\delta A}} \mathcal{F}[A]$

$$\int \mathcal{D}\phi e^{i \int (J + \phi) \mathcal{D}(J + \phi)} \tilde{\mathcal{F}}[\phi] = e^{\frac{i}{2} \int J \mathcal{D} J} e^{-\frac{i}{2} \int \frac{\delta}{\delta A} \mathcal{D} \frac{\delta}{\delta A} \mathcal{F}[A + \int \mathcal{D} J] \Big|_{A \rightarrow 0}}$$

Eikonal Phase (gauge invariant for light-like "velocity" vectors):

$$\int J D J = 2\alpha(0) \cdot \hat{\alpha}(0) \chi(z_T - \bar{z}_T) = 2\alpha(0) \cdot \hat{\alpha}(0) \int_{-\infty}^{\infty} d\tau_1 \int_{-\infty}^{\infty} d\tau_2 n^\mu \bar{n}^\nu \mathcal{D}_{\mu\nu}(z - \bar{z} + \tau_1 n - \tau_2 \bar{n})$$

Collecting all parts:

$$\int \mathcal{D}\alpha \mathcal{D}u \int \mathcal{D}\bar{\alpha} \mathcal{D}\bar{u} e^{i \int (\bar{u} \cdot \bar{\alpha} + u \cdot \alpha)} \left(e^{i \int t \cdot u} \right)_+ \left(e^{i \int t \cdot \bar{u}} \right)_+ e^{i\alpha(0) \cdot \bar{\alpha}(0) \chi(z_T)} e^{-\frac{i}{2} \int \frac{\delta}{\delta A} \mathcal{D} \frac{\delta}{\delta A} \mathcal{F}[A + \int DJ]} \Big|_{A \rightarrow 0}$$

- Quasi-Abelian limit → path-integrals reduce to 1 for $\tau \neq 0$ → ordinary integrals
- Ladder Approximation: Neglect fermion loops and gluon-self interactions: $\mathcal{F} \rightarrow 1$

Final result for the eikonal quark-antiquark scattering amplitude:

$$M_{ab}^{\text{eik}}(q_T) = \frac{2(1-x)P^+}{m_s} \int d^2 z_T e^{-iq_T \cdot z_T} \left[\int d^{N^2-1} \alpha \int \frac{d^{N^2-1} u}{(2\pi)^{N^2-1}} (e^{i\chi(|z_T|)t \cdot \alpha})_{ac} (e^{it \cdot u})_{cb} e^{-i\alpha \cdot u} - 1 \right]$$

Abelian U(1)-theory: recover Coulomb phase

$$\dots \left[e^{i\chi(|z_T|)} - 1 \right]$$

SU(2): analytical evaluation of color effects:

$$e^{i\frac{\sigma}{2} \cdot u} = \cos\left(\frac{|u|}{2}\right) + i \frac{\sigma \cdot u}{|u|} \sin\left(\frac{|u|}{2}\right)$$

SU(3): (?) numerical evaluation, in process...